# Quasi-objective eddy visualization from sparse drifter data 🕫 🚳

Cite as: Chaos **32**, 113143 (2022); https://doi.org/10.1063/5.0099859 Submitted: 19 May 2022 • Accepted: 01 November 2022 • Published Online: 22 November 2022



#### COLLECTIONS

This paper was selected as Featured

SC) This paper was selected as Scilight





## **APL Machine Learning**

Machine Learning for Applied Physics Applied Physics for Machine Learning

Now Open for Submissions



Chaos **32**, 113143 (2022); https://doi.org/10.1063/5.0099859 © 2022 Author(s).

## Quasi-objective eddy visualization from sparse drifter data <a> </a>

Cite as: Chaos **32**, 113143 (2022); doi: 10.1063/5.0099859 Submitted: 19 May 2022 · Accepted: 1 November 2022 · Published Online: 22 November 2022



Alex P. Encinas-Bartos, ២ Nikolas O. Aksamit, and George Haller 🕫 匝

#### AFFILIATIONS

Institute for Mechanical Systems, ETH Zürich, 8092 Zürich, Switzerland

<sup>a)</sup>Author to whom correspondence should be addressed: georgehaller@ethz.ch

#### ABSTRACT

We employ a recently developed single-trajectory Lagrangian diagnostic tool, the trajectory rotation average (TRA), to visualize oceanic vortices (or eddies) from sparse drifter data. We apply the TRA to two drifter data sets that cover various oceanographic scales: the Grand Lagrangian Deployment and the Global Drifter Program. Based on the TRA, we develop a general algorithm that extracts approximate eddy boundaries. We find that the TRA outperforms other available single-trajectory-based eddy detection methodologies on sparse drifter data and identifies eddies on scales that are unresolved by satellite-altimetry.

Published under an exclusive license by AIP Publishing. https://doi.org/10.1063/5.0099859

Meso and submesoscale vortices (or eddies) can trap and transport material over large distances, thereby playing a crucial role in the dynamics of our ecosystem. In order to expand our understanding of the transport of marine tracers, we need to accurately and reliably track the evolution of vortical flow structures. Drifter trajectories represent a valuable but sparse source of information for this purpose. We utilize this information here to evaluate a single-trajectory Lagrangian diagnostic tool that approximates the local material rotation in the flow in a quasi-objective fashion. Our findings on two distinct data sets suggest that this diagnostic accurately highlights material vortices from sparse drifter data.

#### I. INTRODUCTION

Oceanic vortices (or eddies) are highly energetic coherent flow structures that can transport material over large distances. Studying the dynamics of eddies is key to understanding the dispersion of marine tracers, such as biological nutrients and pollutants.<sup>1–5</sup> Lagrangian eddies, generally referred to as elliptic Lagrangian coherent structures (LCSs) in the dynamical systems literature,<sup>6</sup> are material objects that trap and transport floating particles over large distances in the ocean in a coherent fashion. The size of such eddies ranges from a few kilometers (submesoscale) to hundreds of kilometers (mesoscale).

Mesoscale eddies have predominantly been inferred from the sea-surface-height (SSH) field derived from satellite-altimetry data.<sup>7-9</sup> There is, however, increasing evidence that submesoscale currents on the order of a few kilometers influence the marine ecosystem at least as strongly as mesoscale eddies, especially along coastlines and oceanic fronts.<sup>10,11</sup> Despite this observation, submesoscale eddies are rarely studied in detail because their footprint in the SSH field is weak or nonexistent. In contrast to the SSH field, however, drifters follow ocean currents closely and, hence, resolve small-scale features accurately.<sup>12</sup> In addition, surface drifters provide information about the local ocean surface velocity field at a very high temporal resolution. Ocean drifters are, however, sparsely distributed in space, rendering most LCS diagnostics inapplicable to their trajectories. Indeed, most mathematically justifiable algorithms for the detection of elliptic LCSs require differentiation with respect to initial conditions,<sup>6</sup> which is unfeasible for sparse drifter data.

Alternatively, elliptic LCSs can be viewed as coherently evolving sets of material trajectories in space. Based on this view, a wide range of clustering methods for extracting elliptic LCSs have been proposed (see Ref. 13 for a review). These objective methods differ depending on the specific clustering algorithm and the distance metric they employ to define clusters. Common algorithms include fuzzy clustering,<sup>14</sup> spectral graph methods,<sup>15-17</sup> and densitybased clustering methods.<sup>18</sup> Their results, however, rely on user defined input parameters such as the number of clusters, which *a priori* determines the possible elliptic LCS to be detected by the method in the domain. This limitation becomes even more pronounced for sparse drifters for which the number of eddies is *a priori* unknown.

To extract elliptic LCSs from drifter data, one is forced to rely on features of a single trajectory, such as trajectory length, velocity, acceleration, and curvature. These features, however, are all inherently non-objective (frame-dependent) quantities whereas LCSs, as material objects, are objective, i.e., indifferent to coordinate frame changes.<sup>6</sup> Common single-trajectory methods, such as the absolute dispersion,<sup>19</sup> trajectory length,<sup>20,21</sup> Lagrangian spin,<sup>22</sup> maximal trajectory extent,<sup>23</sup> trajectory complexity,<sup>24</sup> and wavelet ridge analysis,<sup>25,26</sup> are limited in capturing elliptic LCSs as they are either non-objective or lack direct physical connection to material deformation in the fluid. Accordingly, while most of these methods were originally introduced to visualize LCSs from truly sparse trajectories, their application to drifter data has remained rare (see Ref. 27 for a review). Notable exceptions include the fully automated looper detection algorithms based on the Lagrangian spin<sup>12,28-30</sup> and the wavelet ridge regression.<sup>31</sup> Both methods seek to extract oscillatory motions from a time-series and lead to comparable results. The methodology employed by the authors in Refs. 12 and 28-30 is an aggregate measure of rotation within a time-series of the Lagrangian spin, whereas Lilly and Pérez-Brunius<sup>31</sup> quantify the instantaneous oscillatory motion of the velocity using signal processing techniques. As a consequence, elliptic LCSs are visualized through looping (or oscillatory) trajectory segments.

In order to distinguish looping from non-looping trajectory segments, a user-defined threshold is inevitably required. In a practical setting, however, differentiating between looping and nonlooping trajectories proves to be challenging and hence finding an appropriate parameter is far from trivial. A common approach is to manually tune a threshold value for looping based on a subset of trajectories. Although this provides a natural and intuitive way to characterize elliptic LCSs, a considerable amount of information is lost when one discards non-looping trajectory segments. It would be more desirable to retain all trajectory information and associate to each trajectory a Lagrangian diagnostic related to material rotation in the flow. This rotational diagnostic could then be plotted over evolving trajectory positions. This approach would provide a qualitative overview of individual vortical flow structures, without relying on any chosen threshold.

Here, we propose to identify eddies from sparse drifter data using the recently introduced adiabatically quasi-objective singletrajectory diagnostics in Ref. 32. These diagnostics closely approximate appropriate objective LCS diagnostics in frames satisfying conditions that typically hold in geophysical flows. We show that the adiabatically quasi-objective trajectory rotation average (TRA) reveals elliptic LCSs (material eddies) at meso- and submesoscales from sparse drifter trajectory data. We additionally compare the vortical flow features extracted from the drifter-based TRA computation with those obtained from other available Lagrangian single-trajectory diagnostics such as the trajectory length<sup>20,21</sup> and the Lagrangian spin.<sup>12,22</sup> We further validate the extracted features with respect to Lagrangian averaged vorticity deviation (LAVD)<sup>33</sup> computed from geostrophic ocean velocity fields derived from satellite-altimetry data (AVISO).

#### II. DATA

## A. Satellite-altimetry ocean-surface current product (AVISO)

The two-dimensional, satellite-altimetry-derived ocean-surface current product (AVISO) has been the focus of several coherent structure studies.<sup>2,34,35</sup> Using this product, the geostrophic velocity  $\mathbf{v}_g(\mathbf{x}, t)$  of ocean currents is obtained from the remotely sensed sea-surface height  $\eta$  as

$$\mathbf{v}_{g}(\mathbf{x},t) = \frac{g}{f} \begin{pmatrix} 0 & -1\\ 1 & 0 \end{pmatrix} \nabla \eta(\mathbf{x},t), \tag{1}$$

where g is the constant of gravity and f is the Coriolis parameter. A global, daily-gridded version of the sea-surface height profile with a spatial resolution of  $(0.25^{\circ} \times 0.25^{\circ})$  is freely available from the Copernicus Marine Environment Monitoring Service.<sup>36</sup>

#### **B.** Drifter data sets

While satellite-based altimetry yields mesoscale velocity fields, surface drifter observations provide direct estimates for the local surface velocity field. In order to illustrate the range of applications of the adiabatically quasi-objective single-trajectory rotation measure introduced in Ref. 32, we will focus on two drifter data sets: the Grand Lagrangian Deployment (GLAD) and the Global Drifter Program (GDP).

#### 1. Grand Lagrangian Deployment (GLAD)

We consider the Coastal Dynamics Experiment (CODE) drifters<sup>37</sup> released during the Grand Lagrangian Deployment (GLAD).<sup>34,38,39</sup> In order to study relative dispersion statistics, around 300 drifters were deployed on 20–31 July 2012 in the northern Gulf of Mexico, sampling various submesoscale features over several weeks. The positions of the drifters were reported every 15 min from which we estimate their velocities via finite differencing. In order to highlight important circulation features, we focus on GLAD drifters active from 10 to 17 August, restricting the domain of interest to

$$\{(x, y) \in [89^{\circ}W, 86^{\circ}W] \times [26^{\circ}N, 29^{\circ}N], \quad t \in [222 \text{ doy}, 229 \text{ doy}]\},$$
(2)

where doy is in short for day of year.

#### 2. Global Drifter Program (GDP)

Additionally, we consider the Global Drifter Program (GDP) data set, which contains more than 20 000 drifters released over the past 40 years. At least 1000 of those drifters are now simultaneously active worldwide.<sup>40</sup> These drifters report their positions every 6 h. We specifically focus on a subset of drifters active from 4 September to 4 October 2006 in the Gulf Stream, i.e., in the domain,

$$\{(x, y) \in [72.5^{\circ}W, 65^{\circ}W] \times [32.5^{\circ}N, 40^{\circ}N], t \in [246 \text{ doy}, 276 \text{ doy}]\}.$$
(3)

#### 3. Drifter data preprocessing

As highlighted by the authors in Refs. 41-43, inertial oscillations have little effect on the overall motion of nearby particles as they periodically return to their starting position after one inertial period. The anticyclonic looping arising from inertial oscillations, however, impacts the drifter velocity profile without affecting separation between nearby particles. In order to remove this unwanted effect from our analysis, we apply a 6th-order low-pass Butterworth filter to the drifter trajectories with a cutoff period of  $T_{cut}$  =  $1.5T_{inertial}$ , as suggested by the authors in Refs. 12, 28, and 41. Here, the inertial period is

$$T_{inertial} = \frac{2\pi}{2\omega\sin(y)},\tag{4}$$

where y is the latitudinal position of the drifter and  $\omega = 7.27 \times 10^{-5} \frac{\text{rad}}{\text{s}}$  is the rotation rate of the Earth.

At mid-latitudes, inertial oscillations have time-scales of 1–2 days, whereas the dominant period of submesoscale and mesoscale eddy motion is above 5 days.<sup>38</sup> The characteristic time of submesoscale and mesoscale features, thus, greatly exceeds the inertial period. Hence, the anticyclonic looping arising from high-frequency diurnal inertial oscillations does not influence the relative dispersion of drifters in the submesoscale and mesoscale regimes.<sup>41</sup> In this paper, we target coherent structures whose time-scales are far above the period of inertial oscillations.

The time interval of definition of a finite-time, temporally aperiodic dynamical system inherently determines the range of features one can identify in that system. Submesoscale eddies have characteristic time-scales of around one week, whereas mesoscale structures arise over months. In contrast, coherent structures tied to inertial oscillations have a temporal range of a few hours. Therefore, in the exploration of short term features, inertial oscillations should not be filtered out of the drifter trajectories.

#### **III. METHODS**

The drifter trajectories,  $\mathbf{x}(t)$ , satisfy the differential equation

 $\dot{\mathbf{x}}(t) = \mathbf{v}(\mathbf{x}, t), \qquad \mathbf{x}(t) \in U \subset \mathbb{R}^2, \quad t \in [t_0, t_N], \quad \mathbf{x}(t_0) = \mathbf{x}_0.$  (5)

Here, **x** is the position, *t* is the time, and  $\mathbf{v}(\mathbf{x}(t), t)$  is the oceansurface velocity field implied by the drifter data. While open-ocean mesoscale features are well captured by satellite-altimetry data,<sup>44</sup> the drifter velocity  $\mathbf{v}(\mathbf{x}, t)$  in coastal regions generally differs from the geostrophic velocity component  $\mathbf{v}_g(\mathbf{x}, t)$  computed from AVISO due to coastal influences (e.g., rivers) and windage.<sup>45</sup> Features derived from drifter trajectories, therefore, generally differ from those obtained from ocean geostrophic velocities.

#### A. Trajectory rotation average

As already noted in Sec. I, LCSs of the velocity field are material sets and, hence, their existence and location are objectives, i.e., indifferent to the observer. All traditionally analyzed features of trajectory data (such as length, curvature, velocity, acceleration, and looping) are, in contrast, observer-dependent and, hence, do not allow for a self-consistent identification of LCSs. To this end,<sup>32,46</sup> developed several quasi-objective diagnostic tools for single trajectories that do approximate objective features of trajectories in frames verifying certain conditions. Inspired by the slowly varying nature of geophysical flow data sets, we restrict our discussion to a family of frames related to each other via slowly varying (or adiabatic) Euclidean coordinate transformations,

$$\mathbf{x} = \mathbf{Q}(t)\mathbf{y} + \mathbf{b}(t), \quad |\mathbf{Q}|, |\mathbf{b}| \ll 1.$$
(6)

Under such slowly varying frame changes, the velocity  $\tilde{\mathbf{v}}$  transforms approximately as an objective vector, i.e.,

$$\tilde{\mathbf{v}} = \mathbf{Q}^T (\mathbf{v} - \dot{\mathbf{Q}}^T \mathbf{y} - \dot{\mathbf{b}}) \sim \mathbf{Q}^T \mathbf{v}, \tag{7}$$

where **v** and  $\tilde{\mathbf{v}}$ , respectively, denote the velocity in the original and in the slowly varying frame. We call a diagnostic adiabatically quasi-objective if in all frames related to each other via Eq. (6), the diagnostic approximates the same objective quantity as long as the frames satisfy a set of additional conditions. Those conditions are specific to the quantity of interest [see Eqs. (9) and (10) for examples].

We will apply such a single-trajectory rotation diagnostic here for the first time to actual sparse drifter data. Similarly to Ref. 32, we consider discretized drifter trajectories  $\{\mathbf{x}(t_i)\}_{i=0}^N$  satisfying Eq. (5). As shown in Refs. 32 and 46, the trajectory rotation average,

$$\overline{\text{TRA}}_{t_0}^{t_N}(\mathbf{x}_0) = \frac{1}{t_N - t_0} \sum_{i=0}^{N-1} \cos^{-1} \frac{\langle \dot{\mathbf{x}}(t_i), \dot{\mathbf{x}}(t_{i+1}) \rangle}{|\dot{\mathbf{x}}(t_i)| \dot{\mathbf{x}}(t_{i+1})|},$$
(8)

approximates a time-averaged trajectory rotation relative to the overall rotation in the flow over the time interval  $[t_0, t_N]$ , in a frame of reference satisfying

$$\frac{\partial \mathbf{v}(\mathbf{x}(t), t)}{\partial t} \ll |\ddot{\mathbf{x}}(t)|. \tag{9}$$

Condition (9) expresses the requirement that Lagrangian time-scales dominate Eulerian time scales. This requirement has been numerically verified for the AVISO data set in Ref. 46 and confirmed in several experimental studies on surface drifters.<sup>37,47,48</sup> As a second condition for  $\overline{\text{TRA}}_{t_0}^{t_N}(\mathbf{x}_0)$  to be an adiabatically quasi-objective scalar field (i.e., approximate an objective scalar field in qualifying frames), the Lagrangian acceleration must dominate the angular acceleration of the trajectory induced by the spatial mean vorticity,

$$\left|\overline{\omega}(t) \times \frac{\dot{\mathbf{x}}(t)}{|\dot{\mathbf{x}}(t)|}\right| \ll \left|\frac{d}{dt}\frac{\dot{\mathbf{x}}(t)}{|\dot{\mathbf{x}}(t)|}\right|.$$
 (10)

We associate to the trajectory  $\mathbf{x}(t)$  over the time interval  $[t_0, t_N]$ its corresponding  $\overline{\text{TRA}}_{t_0}^{t_N}(\mathbf{x}_0)$  value, where  $\mathbf{x}_0 = \mathbf{x}(t_0)$ . Reconstructing the  $\overline{\text{TRA}}_{t_0}^{t_N}(\mathbf{x}_0)$ -field from sparse drifter trajectories via scattered interpolation allows visualization of vortical flow structures. The only parameter involved in the reconstruction of the  $\overline{\text{TRA}}_{t_0}^{t_N}(\mathbf{x}_0)$ field is the choice of the scattered interpolation method. As vortices tend to be elliptic geometric objects, we use linear radial basis function interpolation that favors such structures (see Appendix B for further details). We then apply a spatial average filter of size  $(0.25^{\circ} \times 0.25^{\circ})$  in order to suppress noise but still retain smallscale features. The size of this spatial filter coincides with the spatial

#### ARTICLE

resolution of the AVISO data set. Local maxima in the  $\overline{\text{TRA}}_{t_0}^{\text{IN}}(\mathbf{x}_0)$ -field mark centers of high material rotation and, hence, indicate underlying rotational flow features.

#### **B. Trajectory length**

The trajectory length (also commonly referred to as M-function) is a single-trajectory Lagrangian diagnostic that is applicable to sparse sets of fluid trajectories.<sup>20,21</sup> As pointed out by Ruiz-Herrera,<sup>51,52</sup> this diagnostic is non-objective and has no direct relation to material stretching or rotation. It is nevertheless simple to compute because the arc-length of a trajectory  $\mathbf{x}(t)$  over the time interval  $[t_0, t_N]$  starting at  $\mathbf{x}_0$  is simply

$$M_{t_0}^{t_N}(\mathbf{x}_0) = \sum_{i=0}^{N-1} |\dot{\mathbf{x}}(t_i)| |t_{i+1} - t_i|.$$
(11)

In an attempt to visualize elliptic LCSs from a set of sparse drifter data, we reconstruct  $M_{t_0}^{t_N}(\mathbf{x}_0)$  over a broader domain using linear radial basis functions coupled with a spatial average filter of size (0.25° × 0.25°), as done for the  $\overline{TRA}_{t_0}^{t_N}(\mathbf{x}_0)$ . The authors in Refs. 20, 21, and 53 suggest that dynamical regions in the flow are separated by abrupt variations of  $M_{t_0}^{t_N}(\mathbf{x}_0)$ . Specifically, oceanic eddies are suggested to be signaled by sharp local minima of  $M_{t_0}^{t_N}(\mathbf{x}_0)$ .

#### C. Lagrangian spin

Considering looping trajectories as footprints of coherent Lagrangian eddies,<sup>12,28-30</sup> extract looping drifter segments using the Lagrangian spin

$$\Omega(t_i) = \frac{\langle \mathbf{e}_z, \dot{\mathbf{x}}_{\text{eddy}}(t_i) \times \ddot{\mathbf{x}}_{\text{eddy}}(t_i) \rangle}{\frac{1}{2} |\dot{\mathbf{x}}_{\text{eddy}}(t_i)|^2},$$
(12)

first introduced in Ref. 22. The term  $\dot{\mathbf{x}}_{eddy}(t_i)$  denotes the eddy velocity and  $\mathbf{e}_z = (0, 0, 1)^T$ . In Ref. 29, the eddy velocity  $\dot{\mathbf{x}}_{eddy}(t_i)$  was obtained by removing the background large-scale flow  $\mathbf{u}_{avg}$  from the velocity  $\dot{\mathbf{x}}(t_i)$  of the drifters,

$$\dot{\mathbf{x}}_{\text{eddy}}(t_i) = \dot{\mathbf{x}}(t_i) - \mathbf{u}_{\text{avg}}.$$
(13)

However, recent work points out that removing the background mean current  $\mathbf{u}_{avg}$  results in misidentification of looper segments.<sup>12</sup> Hence, we assume that the drifter velocity is dominated by the eddy fluctuating component rather than the large-scale background flow,

$$\dot{\mathbf{x}}_{\text{eddy}}(t_i) \sim \dot{\mathbf{x}}(t_i).$$
 (14)

The Lagrangian spin  $\Omega$  quantifies the rotation of the velocity vector and is physically related to the vorticity. Evaluating Eq. (12) along a Lagrangian particle trajectory provides an overall measure of rotation within a time-series over the time interval  $[t_0, t_N]$ . Looping segments of a trajectory are characterized as intervals between zero crossings of  $\Omega$ . The duration of each segment of sustained positive or negative  $\Omega$  is referred to as persistence. For each segment, we

define the period P as

$$P = \frac{2\pi}{|\Omega^*|},\tag{15}$$

where  $|\Omega^*|$  is the absolute value of the median of  $\Omega$  over the segment. The time-series of the Lagrangian spin parameter is smoothed using a 3-day median filter in order to suppress noisy observations, as suggested by Lumpkin.<sup>12</sup> Looping segments are defined as trajectory intervals where the persistence exceeds the *ad hoc* chosen threshold value of 2*P*, as proposed in Ref. 12. Decreasing this parameter might reveal additional looping segments, but may also lead to false positives in the identification of loopers. Hence, the identified loopers inherently depend on the choice of the minimum looping period.

#### D. Lagrangian averaged vorticity deviation

Similar to the TRA, the Lagrangian averaged vorticity deviation (LAVD) characterizes the local material rotation in the flow relative to the overall rotation induced by the spatial mean of the vorticity.<sup>33</sup> Computing the LAVD value for a trajectory  $\mathbf{x}(t; \mathbf{x}_0, t_0)$  over the time interval [ $t_0, t_N$ ], from the vorticity

$$\omega(\mathbf{x}(t), t) = \nabla \times \mathbf{v}(\mathbf{x}(t), t)$$

gives

$$LAVD_{t_0}^{t_N}(\mathbf{x}_0) = \frac{1}{t_N - t_0} \int_{t_0}^{t_N} |\omega(\mathbf{x}(s), s) - \overline{\omega}(s)| ds, \qquad (16)$$

where  $\overline{\omega}(t)$  denotes the spatially averaged vorticity at time *t*. Therefore, even though the LAVD is a quantity associated with a single trajectory, its computation requires the knowledge of the velocity field over a large enough domain. This renders the LAVD diagnostics inapplicable to sparse drifter data, even though it has been used to visualize and extract vortices from gridded ocean velocity data sets.<sup>7,33,54</sup> Local maxima in the LAVD-field surrounded by nested, closed, and nearly convex level curves highlight elliptic LCSs. In this work, we apply the LAVD diagnostic to the AVISO data and compare the eddies obtained in this way with those revealed by the TRA from drifter data.

#### E. Validation of eddy boundary extraction algorithm on numerical ocean model

We first validate the eddy boundary extraction algorithm proposed in Appendix A on a numerical ocean model obtained from the AVISO surface velocity field.<sup>36</sup> In our text, we focus on the Agulhas region, which contains a variety of mesoscale eddies analyzed previously by several coherent structure studies.<sup>32,50</sup> We specifically consider the area of the Agulhas leakage,

$$U = \{ (x, y) \in [1, 6] \times [-35, -30] \},$$
(17)

over a period of 25 days. We first compute the LAVD from 200  $\times$  200 densely gridded trajectories and then successively and randomly subsample this trajectory set for a comparative calculation of the TRA. The LAVD highlights three mesoscale eddies [see Fig. 1(a)], whose boundaries are obtained using the algorithm of Haller<sup>50</sup> with the convexity deficiency set to 10<sup>-3</sup>. These three eddies



FIG. 1. LAVD and  $\overline{\text{TRA}}$  evaluated on the full and progressively subsampled AVISO data sets in the Agulhas region. (a) LAVD-field with full resolution (200 × 200). The red curves denote the boundaries of the three LAVD-eddies. [(b)–(d)] Reconstructed  $\overline{\text{TRA}}$ -field from 2000, 200, and 20 randomly selected trajectories. The white curves indicate boundaries of the  $\overline{\text{TRA}}$ -eddies. The black triangles indicate the final trajectory positions.

are clearly visible in the TRA-field [Figs. 1(b) and 1(c)], even after a progressive subsampling of the trajectory density to  $\frac{200}{5^{\circ} \times 5^{\circ}}$ . Upon a further decrease of the trajectory density to  $\frac{20}{5^{\circ} \times 5^{\circ}}$ , two out

of the three mesoscale features still persist in the  $\overline{TRA}$ -field [see Fig. 1(d)]. The third mesoscale feature is not captured by the  $\overline{TRA}$ -field only because no remaining trajectory samples that region after the random sparsification we applied to the original trajectory data set.



FIG. 2. Averaged DICE coefficient plotted as a function of the averaged trajectory spacing per eddy diameter (a) and trajectory density (b). The error bars indicate the minimum and maximum DICE coefficient attained within the corresponding set of trajectories.

We now quantitatively compare the TRA-eddies with the LAVD-eddies using the DICE coefficient that measures the area overlap between the corresponding eddies.<sup>55</sup> We denote the domain covered by the TRA-eddies by Area(TRA<sub>N,eddy</sub>), where *N* is the number of trajectories used to reconstruct the TRA-field. LAVD-eddies that contain at least one trajectory are taken as groundtruth for eddy detection with their domain denoted by Area(LAVD<sub>eddy</sub>). The DICE coefficient for eddy detection comparison is then defined as

$$DICE_{N} = \frac{2(Area(\overline{TRA}_{N,eddy}) \cap Area(LAVD_{eddy}))}{Area(\overline{TRA}_{N,eddy}) + Area(LAVD_{eddy})}.$$
 (18)

A DICE coefficient of 1 would indicate perfect overlap whereas 0 would indicate no overlap between LAVD-eddies and  $\overline{\text{TRA}}$ -eddies.

We vary the number of trajectories from 10 to 2000 and randomly subsample multiple times in order to obtain a statistical estimate of the DICE coefficient for each trajectory density. In Fig. 2, we plot the averaged DICE coefficient and its error bar as a function of two quantities: the averaged trajectory spacing normalized by the eddy diameter [Fig. 2(a)] and the trajectory density [Fig. 2(b)]. The eddy diameter of the three LAVD-eddies in the Agulhas region is around 100 km ( $\sim 1^{\circ}$ ). The averaged trajectory spacing per eddy diameter relates the averaged drifter spacing to the eddy length scales we seek to resolve. Decreasing the drifter density (or, equivalently, increasing the averaged spacing) leads to a drop in the DICE average spacing coefficient. Even for very sparse data sets (e.g., eddy diameter  $\sim$  1), it is possible to approximately identify coherent structures with maximum DICE coefficients of around 0.6. Note, however, that the results inevitably depend on the drifter distribution. If no drifters are inside an eddy, then we do not expect the eddy to be visible in the TRA-field.

GLAD drifters (to be analyzed in Sec. IV A) have an average spacing of 5 km and a trajectory density of around  $\frac{10}{\text{deg} \times \text{deg}}$ . The diameter of the smallest submesoscale eddies observed during the GLAD experiment is around 25 km and the averaged drifter spacing per eddy diameter is 0.2. GDP drifters (to be analyzed in Sec. IV B) are on average separated by 150 km with an estimated trajectory density of  $\frac{0.2}{\text{deg} \times \text{deg}}$ . Mesoscale eddies have characteristic widths ranging from 100 to 300 km<sup>12</sup> and, hence, the averaged spacing per eddy diameter ranges from 0.5 to 1.5. Based on the statistical estimates of the DICE coefficient in Fig. 2, we expect to detect submesoscale and mesoscale eddies in the GLAD and GDP data sets with reasonable confidence.

#### **IV. RESULTS**

#### A. Grand Lagrangian deployment (GLAD)

Drifters released in the GLAD experiment have mostly been used to study dispersion in the ocean over a range of scales.<sup>38,39,41</sup> Furthermore, the authors in Refs. 34 and 35 found a correlation between the evolution of the drifters and nearby attracting LCSs extracted from the geostrophic velocity field. Specifically, drifter behavior agreed with the *tiger tail* pattern inferred from the chlorophyll distribution shown in Fig. 3. A chlorophyll plume extended over more than 100 km from the outlet of the Mississippi River into the open sea and coincided with an attracting LCS.<sup>34</sup> The attracting LCS (continuous black line in Fig. 3) is computed from the geostrophic velocity field  $v_g$  using backward trajectories over the time interval [222 doy, 229 doy] according to Ref. 56. Drifters in the proximity of the tip of the tiger tail organized themselves into long filaments along the attracting LCS. Some of the drifters (in red)



**FIG. 3.** Evolution of GLAD-drifters overlayed on the 8-day composite chlorophyll concentration (12 August 2012). Most drifters (white triangles) formed a long filament along the chlorophyll plume (attracting LCS) extending deep into the Gulf of Mexico. Drifters additionally forming a visible tight cluster are colored red. The blue cross denotes the outlet of the Mississippi River. The black line corresponds to the attracting LCS. (a) t = 222 doy. (b) t = 224 doy. (c) t = 226 doy. (d) t = 229 doy.

additionally exhibited some clustering, suggesting the presence of an elliptic LCS close to the chlorophyll front.

Here, we seek to visualize vortices by reconstructing the  $\overline{\text{TRA}}$ -field from drifter data over the interval [222 doy, 229 doy]

using linear radial basis function interpolation. The TRA, shown in Fig. 4, is plotted with respect to the final drifter positions at time t = 229 doy. This plot reveals multiple rotational features marked by local maxima of the TRA surrounded by a dense



**FIG. 4.** Different material eddy diagnostics computed from the GLAD data set or the AVISO data set. The dashed black lines correspond to the isobaths (-1000, -1500, -2000 m) and highlight the coastal area. (a) Reconstructed  $\overline{\text{TRA}}_{222}^{229}$ -field plotted with respect to the position of the drifters at time t = 229 doy. White crosses indicate local maxima. (b) LAVD $_{222}^{229}$ -field computed from AVISO plotted at time t = 229 doy. The closed red curve denotes the eddy boundary extracted from the LAVD $_{222}^{229}$ -field plotted with respect to the position of the drifters at time t = 229 doy. Red crosses indicate local minima. (d) Cyclonic (red) and anticyclonic (blue) loopers.

set of closed and almost convex contours. For comparison, we include three further Lagrangian eddy diagnostics: the LAVD computed from geostrophic velocity currents, the trajectory length (also called M-function), and the Lagrangian spin computed from drifter data.

#### 1. Coastal area

We start by focusing on the area at the outlet of the Mississippi River on the continental shelf. Coastal flows are regions of intense mixing,<sup>45</sup> often characterized by high potential vorticity and strong



**FIG. 5.** Quasi-materially advected  $\overline{\text{TRA}}_{222}^{229}$  for the GLAD data set over  $t \in [222 \text{ doy}, 229 \text{ doy}] [(a)–(h)]$ . The black line denotes the AVISO-based attracting LCS. Closed white curves mark eddy boundaries extracted from the drifter-based TRA using the algorithm described in Appendix A. Closed red curves indicate eddy boundaries obtained from the AVISO-based LAVD computation. Triangles indicate the positions of the drifters.



**FIG. 6.**  $\overline{\text{KE}}$  evaluated both on GLAD-drifters and on AVISO. The white and red eddies, respectively, indicate the eddies extracted from the drifter-based TRA-field and the AVISO-based LAVD-field. The dashed black lines correspond to the isobaths (-1000, -1500, -2000 m). The black line denotes the AVISO-based attracting LCS. (a) Drifter-based  $\overline{\text{KE}}_{222}^{229}$ -field at time t = 229 doy. White, dashed curves denote a subset of GLAD drifter trajectories contained currently in the same TRA-eddy. (b) AVISO-based  $\overline{\text{KE}}_{222}^{229}$ -field at time t = 229 doy. Red dashed curves denote a subset of AVISO trajectories contained currently in the same TRA-eddy.

horizontal shear, which are responsible for the formation of smallscale eddies.<sup>57,58</sup> Anticyclonic looping segments extracted from the Lagrangian spin parameter confirm the presence of small-scale vortices at the outlet of the Mississippi River [Fig. 4(d)]. Evidence for the existence of submesoscale elliptic features and strong mixing areas on the continental shelf is also found in the  $\overline{\text{TRA}}$ -field [Fig. 4(a)]. Vortices are revealed in the  $\overline{\text{TRA}}$  as local maxima surrounded by a dense set of closed curves, indicating abrupt spatial variations. The local maximum of the  $\overline{\text{TRA}}$  at (88.6°W, 28.1°N) is surrounded by a dense set of closed contours, thereby indicating high spatial gradients in the  $\overline{\text{TRA}}$ -field. Close to the vortical flow features identified by the TRA in the coastal area, the M-function displays two local minima, respectively, at (88.6°W, 28.2°N) and (88.5°W,27.8°N) [Fig. 4(c)]. Similarly to the TRA and the Lagrangian spin parameter, the M-function also indicates the existence of coastal eddies. However, compared to the TRA, which displays a unique local maximum surrounded by a dense set of closed and convex curves, the M-function displays multiple local minima surrounded by sparse level sets, thereby indicating a region without sharp minima.

The LAVD-field displays no distinguished features close to the outlet of the Mississippi River [Fig. 4(b)]. Eddy boundaries can be selected as outermost (almost) convex contours around a local maximum in the LAVD-field.<sup>33</sup> Passing from the LAVD-field to a discrete

set of closed curves requires specifying two parameters: the minimum length  $l_{min}$  of the perimeter of the eddy and the convexity deficiency  $c_d$ . The convexity deficiency is

$$c_{\rm d} = \frac{|A_{\rm contour} - A_{\rm contour, convex}|}{A_{\rm contour}},$$
(19)

where  $A_{contour}$  is the area enclosed by the closed contour and  $A_{contour,convex}$  is the area enclosed by the convex hull of the closed contour. Similarly to Ref. 33, we set  $c_d = 10^{-3}$ . As a threshold for the perimeter of the eddy  $l_{min}$ , we select  $l_{min} = 2\pi r$ , with r = 25 km (~ 0.25°). This corresponds to the minimal length scales that can reliably be resolved by the AVISO data set with a spatial resolution of  $(0.25^{\circ} \times 0.25^{\circ})$ . The LAVD does not reveal the presence of coastal eddies [Fig. 4(b)]. We attribute the mismatch between the LAVD and the above discussed drifter-based diagnostics to the insufficient resolution of the underlying ocean surface velocity close to the outlet of the Mississippi River by the AVISO data set. Hence, in coastal areas, the drifter-based TRA diagnostic captures the flow dynamics with higher detail than the AVISO-based LAVD.

#### 2. Open sea

The TRA highlights another family of elliptic LCSs close to the chlorophyll plume extending from the outlet of the Mississippi



**FIG. 7.** Lagrangian diagnostics evaluated on a subset of the GDP data set. (a) Reconstructed  $\overline{TRA}_{246}^{276}$ -field plotted with respect to the position of the drifters (triangles) at time t = 276 doy. White cross indicates a unique local maximum in the  $\overline{TRA}_{246}^{276}$ -field. (b) Eddy boundary superimposed on the floating sargassum concentration inferred from the Maximum Chlorophyll Index (MCI) at t = 276 doy. (c) Reconstructed  $M_{246}^{276}$ -field plotted with respect to the position of the drifters (triangles) at time t = 276 doy. White cross indicates a unique local maximum in the  $M_{246}^{276}$ -field plotted with respect to the position of the drifters (triangles) at time t = 276 doy. White cross indicates a unique local maximum in the  $M_{246}^{276}$ -field plotted with respect to the position of the drifters (triangles) at time t = 276 doy. White cross indicates a unique local maximum in the  $M_{246}^{276}$ -field. (d) Cyclonic (red) loopers.

River to the open ocean [Fig. 4(a)]. The two local maxima marking elliptic LCSs are located at (87.5°W, 26.8°N) and (87.3°W, 27.4°N). Due to the close proximity of these extrema, it is, however, *a priori* unclear whether they highlight two separate eddies or whether they

are part of the same mesoscale eddy. The cyclonic looping segments confirm the presence of vortical flow features in this area [red trajectories in Fig. 4(d)], but do not specifically highlight eddy boundaries. Inspection of the cyclonic trajectories suggests that they are

originally part of two distinct eddies as they come from two separate regions. Extracting looping trajectory segments from the time-series of the Lagrangian spin is a powerful methodology. However, discriminating between loopers and non-loopers requires an *ad hoc* choice of the minimum sustained looping period of a trajectory. In contrast, the TRA retains all trajectory information and allows visualization of vortical flow structures from sparse drifter data using a scalar diagnostic.

The investigated elliptic LCSs highlighted by the  $\overline{\text{TRA}}$  and the Lagrangian spin are also visible in the LAVD, which displays a nearly convex vortical flow feature [closed red curve in Fig. 4(b)]. Compared to the  $\overline{\text{TRA}}$ , however, which indicates two separate albeit closely located local maxima, the LAVD computation clearly reveals a single mesoscale eddy.

In contrast to the aforementioned methods, the features resulting from the trajectory length diagnostic are far less pronounced [Fig. 4(c)]. Indeed, the M-function displays multiple local minima, which are only partially correlated with the eddies suggested by the other methods. Furthermore, local minima in the M-function are not sharp, as indicated by the sparsity of the surrounding level curves. Hence, extracting eddy-related features from the M-function is challenging as there exist no distinguishable sharp and closed boundaries surrounding the local minima.

#### 3. Eddy dynamics

In order to investigate the formation and evolution of the elliptic LCSs inferred from the drifter-based  $\overline{\text{TRA}}$ , we proceed by quasi-materially advecting the  $\overline{\text{TRA}}$  distribution over the time interval [222 doy, 229 doy] (Fig. 5). True material advection would require a spatiotemporally well-resolved velocity field. In our setting, however, the velocity field is only sparsely known and, hence, the advected structures are inherently non-material: At every time step, the  $\overline{\text{TRA}}$ -field must be approximated from the current drifter distribution. As a consequence, the evolution of the extracted eddy boundaries is not exactly Lagrangian. This implies that a set of drifters may not stay inside an eddy over the full time interval. For the same reason, the extracted eddies can potentially merge.

The closed white curves in Fig. 5 indicate eddy boundaries extracted from the TRA using the algorithm described in Appendix A. The red closed curve denotes the truly materially advected vortex boundary extracted from the LAVD at time t = 229 doy. The LAVD-based eddy at t = 229 doy is materially advected using the geostrophic velocity  $\mathbf{v}_g(\mathbf{x}, t)$ , whereas the eddy boundaries inferred from the drifter-based TRA are quasi-materially advected along drifter trajectories.

At t = 222 doy, the TRA suggests the presence of several smallscale vortices at the outlet of the Mississippi River and at open sea [Fig. 5(a)]. The submesoscale eddies close to the outlet of the Mississippi River remain trapped in coastal areas and eventually merge into a larger vortical flow feature. Over the time interval [222 doy, 226 doy], the LAVD-based eddy does not coincide with any of the eddies inferred from the TRA [Figs. 5(a)–5(d)]. The white eddy initially located at approximately (88.5°W, 27.5°N) is associated with the clustered red drifters identified in Fig. 3, thereby confirming the existence of the submesoscale eddy along the chlorophyll front, which agrees with the observation put forward in Refs. 35 and 34. The eddy develops along the attracting LCS and then slowly detaches away from the oceanic front [Figs. 5(a)-5(f)]. Drifters are thereby coherently transported from the coastline into the open sea. The drifter-based eddy evolving along the AVISObased attracting LCS eventually merges with the submesoscale eddy originally located at (86.5°W, 27.25°N) to form a larger mesoscale eddy at t = 228 doy [Fig. 5(g)]. Hence, elliptic LCSs generated along oceanic fronts offer a transport mechanism for particles, carrying material over long distances away from the coastline into the open sea. We point out that intersections between attracting and elliptic LCS (such as at t = 222 doy) are physically inconsistent as they imply contradicting material response. We attribute this inconsistency to the fact that the attracting LCS is computed from  $\mathbf{v}_{\alpha}(\mathbf{x}, t)$  whereas the white elliptic LCSs are computed from drifter velocities.

Toward the end of the advection process, the LAVD-based elliptic LCSs [red curve in Fig. 5(h)] approximates the mesoscale eddy inferred from the TRA [white curve centered at approximately  $(87.5^{\circ}W, 27^{\circ}N)$  in Fig. 5(h)]. The red eddy shows no degree of filamentation and remains coherent over the full time interval. The white eddy results from the vortex merger between two smaller eddies and is significantly larger than the red eddy. At t = 229 doy, the circumference of the TRA-based white eddy and the LAVD-based red eddy is 298.5 and 152 km. As the LAVDbased eddy is a materially advected closed curve, it can neither split nor merge with any other materially advected curve. Hence, by construction, the LAVD is unable to capture vortex mergers. In contrast, the quasi-materially advected, TRA-based eddies can merge into larger eddies. This follows because the evolving eddies are not purely Lagrangian, given that the computation of the eddy boundary from the TRA-field is independently carried out at each time step.

Kinetic energy (KE) is frequently seen as an indicator of largescale coherent motion in the ocean.<sup>12,59,60</sup> The averaged KE associated with a trajectory  $\mathbf{x}(t)$  over the time interval  $[t_0, t_N]$  starting at  $\mathbf{x}_0$ is

$$\overline{\mathrm{KE}}_{t_0}^{\mathbf{t}_N}(\mathbf{x}_0) = \frac{1}{2|t_N - t_0|} \sum_{i=0}^{N-1} |\dot{\mathbf{x}}(t_i)|^2 |t_{i+1} - t_i|, \qquad (20)$$

where  $\overline{\text{KE}}$  is a non-objective, Lagrangian single-trajectory diagnostics that bears similarities to the trajectory length diagnostic introduced in Sec. III B. It is often used to visualize energetically dominant flow structures both from AVISO<sup>61</sup> and drifter data.<sup>12</sup> Here, we compute the AVISO-based  $\overline{\text{KE}}$ -field from synthetic Lagrangian particle trajectories generated by the geostrophic velocity field  $\mathbf{v}_g(\mathbf{x}, t)$ in the region of interest [Fig. 6(b)]. The computed  $\overline{\text{KE}}$ -field displays a front-like feature coinciding with the attracting LCS responsible for the transport drifters from the outlet of the Mississippi River into the open sea. The drifter-based  $\overline{\text{KE}}$ -field reconstructed from drifter data shows a front-like feature reminiscent of the front visible in the AVISO-based  $\overline{\text{KE}}$  [Fig. 6(a)]. The weakly spiraling feature in the AVISO-based  $\overline{\text{KE}}$ -field corresponds to the mesoscale elliptic LCSs. Overall, however, both methodologies fail to clearly highlight the oceanic eddies detected from TRA and LAVD. Hence, although frequently related to coherent eddy motions, we conclude that the kinetic energy is not conclusively linked to oceanic eddies in the region of interest. Figure 6 also shows drifters (white) and AVISO trajectories (red) released within TRA-based eddies. Overall, the real and synthetic drifter trajectories divert from each other over time, as expected. The difference is more pronounced in coastal areas due to the influence of external factors such as the outflow of the Mississippi-River.

The quasi-material advection of the  $\overline{\text{TRA}}$ -based eddies sheds new light on the origin and the formation of the mesoscale eddy detected from the AVISO-based LAVD-field. Despite the minimal amount of data, the  $\overline{\text{TRA}}$  reveals eddy dynamics hidden in the LAVD and  $\overline{\text{KE}}$ .

#### B. Global Drifter Program (GDP)

In our second example, we focus on a set of drifters in the western North-Atlantic. This oceanic region is characterized by a strong and persistent formation of eddies arising from the meanders of the Gulf Stream.<sup>62,63</sup> On 4 October 2006, a floating sargassum patch was detected by the Medium Resolution Imaging Spectrometer (MERIS) on Envisat. This feature has a spiralling shape that is also visible from satellite-altimetry data.<sup>2</sup> This floating sargassum patch is visualized in Fig. 7(b) using the Maximum Chlorophyll Index (MCI).<sup>64</sup> Due to frequent cloud coverage, such clear snapshots of floating material in the ocean are very rare.

We use the  $\overline{\text{TRA}}$  to extract the eddy highlighted by the spiralling sargassum patch described in Ref. 2. We also compute two further single trajectory metrics: the trajectory length diagnostic [Fig. 7(c)] and the looping segments derived from the Lagrangian spin [Fig. 7(d)]. We include a snapshot of the floating sargassum patch that reveals the presence of an elliptic LCS [Fig. 7(b)]. Similarly to the MCI, the TRA reveals a Lagrangian eddy centered at (67.6°W, 37.1°N). Indeed, this eddy is visible as a distinguished local maximum in the TRA-field surrounded by a dense set of closed and convex contours [the zoomed inset of Fig. 7(a)]. The white eddy boundary is extracted from the  $\overline{\text{TRA}}$  using the algorithm proposed in Appendix A with the same parameters presented in Sec. IV A 3. The extracted eddy boundary underestimates the size of the sargassum patch, but correctly approximates the location [Fig. 7(b)]. The cyclonic looping exhibited by the two trajectories inside the eddy additionally confirms the presence of an elliptic LCS [Fig. 7(d)] but does not provide a specific estimate for the eddy boundary. The trajectory length diagnostic shows a nearby maximum [the zoomed inset of Fig. 7(c)], which is inconsistent with the generally suggested footprint (local minima) for a coherent eddy in the M-field (see Sec. III B).

Figure 8 shows the LAVD-field with AVISO trajectories (red) launched backwards in time from true drifter positions at time t = 276 doy. The mesoscale eddy inferred from the sparse drifter distribution (white curve in the zoomed inset of Fig. 8) is in close agreement with the LAVD-eddy (red curve in the zoomed inset of Fig. 8). As expected, drifter and AVISO trajectories show similar looping behavior inside the identified coherent structure, while they largely differ in other regions.



FIG. 8. AVISO-based trajectories (red) and true drifter trajectories (white) superimposed on LAVD-field. The closed white and red curves in the zoomed inset, respectively, indicate the TRA-and LAVD-eddy. Triangles denote drifter positions.

#### **V. CONCLUSION**

Lagrangian eddies (elliptic LCSs) are material objects responsible for the transport of floating particles over large distances in the ocean. They are, by their definition, frame-indifferent and, thus, can only be reliably deduced from objective feature extraction methods. The local ocean velocity is most accurately observed from float trajectory data, which, however, is inherently non-objective, representing an inconsistency that available eddy detection methods for sparse trajectories do not address. Those methods typically describe eddies by extracting the looping segments of a trajectory, but their definition of looping depends on the frame of the observer. Furthermore, looping segments of a trajectory are most commonly described by these methods in a statistical sense and, hence, are not geared toward highlighting individual Lagrangian eddy boundaries with high accuracy.

In this paper, we have proposed to tackle this inconsistency from a dynamical systems perspective by applying the adiabatically quasi-objective TRA diagnostic<sup>32,46</sup> to sparse drifter data sets. The TRA approximates an objective measure of material rotation in frames satisfying specific conditions that generally hold in the ocean. We have found that vortical flow features are related to regions of high local material rotation identified as local maxima in the TRAfield. The TRA highlights both submesoscale and mesoscale vortices from sparse drifter data, as demonstrated in our two examples. Furthermore, it also succeeds in characterizing the mixing and stirring processes in coastal flow regions and captures the merger of two originally distinct eddies. Contrary to other single trajectory diagnostics, both the TRA and the LAVD are physically related to the local material rotation in the flow. In contrast to the LAVD, which correctly highlights vortical flow features in sufficiently dense velocity data, the TRA can be applied to arbitrarily sparse drifter data, given its lack of dependence on nearby drifter trajectories. Hence, it incorporates valuable drifter data into the analysis of oceanic coherent structures in a physically and mathematically justifiable way. This proves to be especially useful in ocean regions where satellite-altimetry data do not unravel the true ocean dynamics.

Importantly, the looper segments extracted from the Lagrangian spin coincide with the features identified in the  $\overline{TRA}$ . However, a spaghetti plot of looping trajectory segments does not immediately reveal transport barriers and eddy boundaries. Furthermore, potentially valuable information is lost when we discard non-looping trajectory segments based on a manually tuned threshold parameter. The trajectory length diagnostic<sup>20,21</sup> also tended to show either minima or maxima near the material eddies highlighted by the local maxima of the  $\overline{TRA}$ -field. This variation in extremum types creates ambiguity in using the trajectory length as a stand-alone indicator for detecting elliptic LCSs from sparse data sets.

Apart from the visual inspection of the reconstructed TRAfield, we have additionally presented an algorithm to extract approximate eddy boundaries from sparse drifter data. As vortical flow features are indicated by blobs close to local maxima in the TRAfield, the proposed method resembles a blob detection algorithm. Passing from a continuous scalar diagnostic field to a set of closed curves inevitably requires introducing user-defined parameters. All in all, however, the number of free parameters is comparable to other multi-trajectory Lagrangian eddy detection methods.<sup>33,65</sup> This is noteworthy as these algorithms were originally designed assuming knowledge of the underlying velocity field.

#### ACKNOWLEDGMENTS

The authors acknowledge financial support from Priority Program SPP 1881 (Turbulent Superstructures) of the German National Science Foundation (DFG).

#### AUTHOR DECLARATIONS

#### **Conflict of Interest**

The authors have no conflicts to disclose.

#### **Author Contributions**

Alex P. Encinas Bartos: Data curation (equal); Investigation (equal); Writing - original draft (equal). Nikolas O. Aksamit: Supervision (equal); Writing - review & editing (equal). George Haller: Conceptualization (equal); Funding acquisition (equal); Supervision (equal); Writing - review & editing (equal).

#### DATA AVAILABILITY

The AVISO geostrophic current velocity product used in this study "Global Ocean Gridded L4 Sea Surface Heights and Derived Variables Reprocessed" is freely available and is hosted by the Copernicus Marine Environment Monitoring Service.<sup>36</sup> The GDP

drifter data is openly available in the NOAA Global Drifter Program data set.66 The GLAD drifter data is distributed by the Gulf of Mexico Research Initiative Information and Data Cooperative (GRIIDC).67 The attracting LCS and LAVD computations have been performed with the software package TBarrier. Jupyter notebooks (together with drifter data) implementing the methods used here are available under Github/EncinasBartos. These notebooks can readily be applied to any sparse trajectory data set.

#### APPENDIX A: EDDY BOUNDARY EXTRACTION ALGORITHM

While various eddy extraction algorithms have been presented in Refs. 16, 17, 50, and 68, these methods assume a trajectory density that is generally unavailable for drifter trajectories in the ocean. Here, we propose an algorithm that extracts approximate eddy boundaries from the topology of the reconstructed TRA-field. Vortices are then identified by this algorithm as local maxima of the TRA-field surrounded by a dense set of closed and convex curves characterized by high spatial gradients.

Passing from a continuous scalar field to a set of discrete closed curves representing eddy boundaries inevitably requires introducing threshold parameters. There are two main parameters involved in Algorithm 1. The first user-defined quantity aids the identification of the local maxima in the  $\overline{TRA}$ -field. As we identify vortical flow features with regions of high  $\overline{\text{TRA}}$ , local maxima below a predefined threshold  $\overline{\text{TRA}}_{\text{loc,max}}$  are neglected. Additionally, we also need to specify the minimum number of drifters  $n_d$  inside an eddy. As elliptic LCSs are often observed via a dense clustering of multiple drifters, this parameter is generally set to be greater than 1. In this

Algorithm 1. Extraction of approximate eddy boundaries from  $\overline{TRA}_{t_n}^{t_N}$ -field

**Input:** Trajectories over the time interval  $[t_0, t_N]$ .

- 1. Reconstruct  $\overline{\text{TRA}}_{t_0}^{t_N}$ -field at time *t* using linear radial basis interpolation. It is recommended to additionally filter the resulting  $\overline{\text{TRA}}_{t_0}^{t_N}$  using a spatial average filter to reduce noise. 2. Find local maxima of  $\overline{\text{TRA}}_{t_0}^{t_N}$  which are above a threshold
- TRA<sub>loc,max</sub>.
- 3. Compute for each closed level set surrounding a local maximum, the averaged  $|\nabla \overline{\text{TRA}}_{t_0}^{t_N}|$  along the level set.
- 4. Find closed level set with the highest averaged  $|\nabla \overline{TRA}_{t_0}^{t_N}|$  , which additionally

(a) has at least one local maximum of  $\overline{TRA}_{t_0}^{t_N}$  in its interior.

(b) contains at least n<sub>d</sub>-trajectories.

- 5. Take the convex hull of all the selected closed curves.
- 6. If two or more convex closed curves intersect, then take the union of these curves.
- 7. Take the convex hull of the resulting closed curves.
- Output: Approximate eddy boundaries at time t.



**FIG. 9.** Comparison of TRA-fields obtained from different interpolation functions with the raw TRA plot for the GLAD data set at time t = 229 doy. Closed white curves indicate eddies extracted from the reconstructed TRA-field. White crosses, circles, and squares denote local TRA maxima. Triangles indicate the position of the drifters. (a) Linear rbf, (b) linear ( $C^0$ -interpolation), (c) natural neighbor ( $C^1$ -interpolation), and (d) scattered, raw TRA. Color and size of the triangles are related to the TRA-value associated with each drifter.

work, the parameters are consistently set to  $\overline{TRA}_{loc,max} = .5\overline{TRA}_{max}$ , where  $\overline{TRA}_{max}$  is the global maximum in the  $\overline{TRA}$ -field and  $n_d = 2$ , as we expect multiple drifters to be part of the same coherent structure. Hence, our eddy extraction algorithm detects elliptic LCSs if at

least two trajectories are contained inside an eddy. By construction, eddy boundaries are closed convex curves characterized by sharp gradients. These curves do not necessarily coincide with level sets of the  $\overline{\text{TRA}}$ .

## APPENDIX B: SENSITIVITY ANALYSIS WITH RESPECT TO THE INTERPOLATION METHOD

The features of the reconstructed  $\overline{\text{TRA}}$ -field clearly depend on the employed interpolation scheme, but we expect the topology of the  $\overline{\text{TRA}}$ -field to be robust with respect to the interpolation method in the vicinity of sharp  $\overline{\text{TRA}}$  maxima. Here, we verify the persistence of such local maxima and the extracted  $\overline{\text{TRA}}$ -eddies with respect to common interpolants: linear radial basis function (rbf), linear ( $C^0$ -interpolant), and natural neighbor ( $C^1$ -interpolant). The reconstructed scalar fields will all be equally pre-and post-processed: inertial oscillations are removed from the drifter trajectories and



**FIG. 10.** Same as Fig. 9 but for the GDP data set at t = 276 doy.

a spatial averaging filter of size  $(0.25^\circ\times0.25^\circ)$  is applied to the reconstructed  $\overline{TRA}\text{-field}.$ 

Figure 9 shows the reconstructed TRA-field using linear rbf, linear, and natural neighbor interpolation for the GLAD drifters. For comparison, we also included the raw TRA plot. Two major eddies are detected independent of the interpolation scheme [see Figs. 9(a)–9(c)]. Both the linear rbf and natural neighbor interpolation reveal three similar local maxima [white crosses/squares in Figs. 9(a) and 9(c)]. On average, the white crosses and squares are only separated by 1 km (~0.01°) and the average distance of the local maxima to the closest drifter is around 3 km (~0.03°) in both cases.

The linear interpolant reveals three further local maxima closely located to those suggested by the linear rbf and natural neighbor interpolation scheme [white circles in Fig. 9(b)]. These additional local maxima, however, are not strong enough to lead to the emergence of new eddies. Furthermore, local maxima are located in regions with dense TRA-values [see Fig. 9(d)]. All three interpolation schemes suggest the emergence of two robustly persisting eddies.

We perform an analogous sensitivity analysis on the reconstructed TRA-field for the GDP drifters at time t = 276 doy (see Fig. 10). Local maxima of the linear rbf and natural neighbor interpolation are in close agreement with each other, whereas the linear interpolant introduces two further local maxima. Again, these additional local maxima are weak and do not lead to the detection of additional eddies. In summary, the eddy boundary extraction algorithm consistently reveals a mesoscale eddy independent of the interpolation scheme.

Overall, we find that the eddy location does not strongly depend on the interpolation scheme at least in the two data sets considered in this paper. However, the exact shape and size of the eddies vary as a function of the interpolation method. In very sparse data sets, such as the GDP-data set, the identified eddy area may vary substantially under changes in the interpolation method. The location of the largest local TRA maximum signaling the presence of an eddy, however, is fairly robust with respect to the interpolation method. Due to its inherent radial symmetry, linear rbf interpolation tends to favor elliptic structures. In contrast, linear and natural neighbor interpolation leads to sharper and non-smooth eddy boundaries.

#### REFERENCES

<sup>1</sup>R. Abernathey, C. Bladwell, G. Froyland, and K. Sakellariou, "Deep Lagrangian connectivity in the global ocean inferred from Argo floats," J. Phys. Oceanogr. **52**, 951–963 (2021).

<sup>2</sup>F. J. Beron-Vera, M. J. Olascoaga, G. Haller, M. Farazmand, J. Triñanes, and Y. Wang, "Dissipative inertial transport patterns near coherent Lagrangian eddies in the ocean," Chaos **25**, 087412 (2015).

<sup>3</sup>G. Froyland, R. M. Stuart, and E. van Sebille, "How well-connected is the surface of the global ocean?," Chaos **24**, 033126 (2014).

<sup>4</sup>P. Miron, M. J. Olascoaga, F. J. Beron-Vera, N. F. Putman, J. Triñanes, R. Lumpkin, and G. J. Goni, "Clustering of marine debris and sargassum like drifters explained by inertial particle dynamics," Geophys. Res. Lett. **47**, e2020GL089874, https://doi.org/10.1029/2020GL089874 (2020).

<sup>5</sup>E. van Sebille, M. H. England, and G. Froyland, "Origin, dynamics and evolution of ocean garbage patches from observed surface drifters," Environ. Res. Lett. 7, 044040 (2012).

<sup>6</sup>G. Haller, "Lagrangian coherent structures," Annu. Rev. Fluid Mech. 47, 137–162 (2015). <sup>7</sup>R. Abernathey and G. Haller, "Transport by Lagrangian vortices in the eastern Pacific," J. Phys. Oceanogr. **48**(3), 667–685 (2018).

<sup>8</sup>D. B. Chelton, M. G. Schlax, R. M. Samelson, and R. A. de Szoeke, "Global observations of large oceanic eddies," Geophys. Res. Lett. **34**(15), L15606, https://doi.org/10.1029/2007GL030812 (2007).

<sup>9</sup>J. H. Faghmous, I. Frenger, Y. Yao, R. Warmka, A. Lindell, and V. Kumar, "A daily global mesoscale ocean eddy dataset from satellite altimetry," Sci. Data 2, 1–16 (2015).

<sup>10</sup>M. Lévy, P. J. Franks, and K. S. Smith, "The role of submesoscale currents in structuring marine ecosystems," Nat. Commun. 9, 4758 (2018).

<sup>11</sup>J. C. McWilliams, "Submesoscale currents in the ocean," Proc. R. Soc. A **472**, 20160117 (2016).

<sup>12</sup>R. Lumpkin, "Global characteristics of coherent vortices from surface drifter trajectories," J. Geophys. Res.: Oceans **121**, 1306–1321, https://doi.org/10.1002/ 2015JC011435 (2016).

<sup>13</sup>A. Hadjighasem, M. Farazmand, D. Blazevski, G. Froyland, and G. Haller, "A critical comparison of Lagrangian methods for coherent structure detection," Chaos 27(5), 053104 (2017).

<sup>14</sup>G. Froyland and K. Padberg-Gehle, "A rough-and-ready cluster-based approach for extracting finite-time coherent sets from sparse and incomplete trajectory data," Chaos 25, 087406 (2015).

<sup>15</sup>M. Filippi, I. I. Rypina, A. Hadjighasem, and T. Peacock, "An optimizedparameter spectral clustering approach to coherent structure detection in geophysical flows," Fluids **6**(1), 39 (2021).

physical flows," Fluids **6**(1), 59 (2021). <sup>16</sup>A. Hadjighasem, D. Karrasch, H. Teramoto, and G. Haller, "Spectralclustering approach to Lagrangian vortex detection," Phys. Rev. E **93**, 063107 (2016).

<sup>17</sup>K. L. Schlueter-Kuck and J. O. Dabiri, "Coherent structure colouring: Identification of coherent structures from sparse data using graph theory," J. Fluid Mech. 811, 468–486 (2017).

<sup>18</sup>S. Mowlavi, M. Serra, E. Maiorino, and L. Mahadevan, "Detecting Lagrangian coherent structures from sparse and noisy trajectory data," J. Fluid Mech. **948**, A4 (2022).

<sup>19</sup>A. Provenzale, "Transport by coherent barotropic vortices," Annu. Rev. Fluid Mech. **31**, 55–93 (1999).

<sup>20</sup>A. M. Mancho, S. Wiggins, J. Curbelo, and C. Mendoza, "Lagrangian descriptors: A method for revealing phase space structures of general time dependent dynamical systems," Commun. Nonlinear Sci. Numer. Simul. 18, 3530–3557 (2013).

<sup>21</sup>C. Mendoza and A. M. Mancho, "Hidden geometry of ocean flows," Phys. Rev. Lett. **105**(3), 038501 (2010).

<sup>22</sup>B. L. Sawford, "Rotation of trajectories in Lagrangian stochastic models of turbulent dispersion," Boundary-Layer Meteorol. 93, 411–424 (1999).

<sup>23</sup>R. Mundel, E. Fredj, H. Gildor, and V. Rom-Kedar, "New Lagrangian diagnostics for characterizing fluid flow mixing," Phys. Fluids 26, 126602 (2014).

<sup>24</sup>I. I. Rypina, S. E. Scott, L. J. Pratt, and M. G. Brown, "Investigating the connection between complexity of isolated trajectories and Lagrangian coherent structures," Nonlinear Processes Geophys. **18**, 977–987 (2011).

<sup>25</sup>J. M. Lilly and S. C. Olhede, "Bivariate instantaneous frequency and bandwidth," IEEE Trans. Signal Process. 58, 591–603 (2009).

<sup>26</sup>J. M. Lilly and S. C. Olhede, "On the analytic wavelet transform," IEEE Trans. Inf. Theory 56(8), 4135–4156 (2010).

<sup>27</sup>I. I. Rypina, T. Getscher, L. J. Pratt, and T. Ozgokmen, "Applying dynamical systems techniques to real ocean drifters," Nonlinear Processes Geophys. 29(4), 345–361 (2022).

<sup>28</sup>C. Dong, Y. Liu, R. Lumpkin, M. Lankhorst, D. Chen, J. C. McWilliams, and Y. Guan, "A scheme to identify loops from trajectories of oceanic surface drifters: An application in the Kuroshio Extension region," J. Atmos. Oceanic Technol. 28, 1167–1176 (2011).

<sup>29</sup>A. Griffa, R. Lumpkin, and M. Veneziani, "Cyclonic and anticyclonic motion in the upper ocean," Geophys. Res. Lett. **35**, L01608, https://doi.org/10.1029/2007GL 032100 (2008).

<sup>30</sup>M. Veneziani, A. Griffa, A. M. Reynolds, and A. J. Mariano, "Oceanic turbulence and stochastic models from subsurface Lagrangian data for the Northwest Atlantic Ocean," J. Phys. Oceanogr. **34**(8), 1884–1906 (2004). <sup>31</sup> J. M. Lilly and P. Pérez-Brunius, "Extracting statistically significant eddy signals from large Lagrangian datasets using wavelet ridge analysis, with application to the Gulf of Mexico," Nonlinear Processes Geophys., 28, 181-212 (2021).

32G. Haller, N. O. Aksamit, and A. P. Encinas-Bartos, "Quasi-objective coherent structure diagnostics from single trajectories," Chaos 31, 043131 (2021).

<sup>33</sup>G. Haller, A. Hadjighasem, M. Farazmand, and F. Huhn, "Defining coherent vortices objectively from the vorticity," J. Fluid Mech. 795, 136-173 (2016).

34 M. J. Olascoaga, F. J. Beron-Vera, G. Haller, J. Triñanes, M. Iskandarani, E. F. Coelho, B. K. Haus, H. S. Huntley, G. Jacobs, A. D. Kirwan, and B. L. Lipphardt, "Drifter motion in the Gulf of Mexico constrained by altimetric Lagrangian coherent structures," Geophys. Res. Lett. 40, 6171-6175, https://doi.org/10.1002/2013GL058624 (2013).

<sup>35</sup>M. J. Olascoaga and G. Haller, "Forecasting sudden changes in environmental pollution patterns," Proc. Natl. Acad. Sci. 109, 4738–4743 (2012). <sup>36</sup>See https://doi.org/10.48670/moi-00145 for "E.U. Copernicus Marine Service

Information."

<sup>37</sup>R. E. Davis, "Drifter observations of coastal surface currents during CODE: The method and descriptive view," J. Geophys. Res.: Oceans 90, 4741-4755, https://doi.org/10.1029/JC090iC03p04741 (1985).

38 A. J. Mariano, E. H. Ryan, H. S. Huntley, L. C. Laurindo, E. Coelho, A. Griffa, and M. Wei, "Statistical properties of the surface velocity field in the northern Gulf of Mexico sampled by GLAD drifters," J. Geophys. Res.: Oceans 121, 5193-5216, https://doi.org/10.1002/2015JC011569 (2016).

<sup>39</sup>A. C. Poje, T. M. Özgökmen, B. L. Lipphardt, B. K. Haus, E. H. Ryan, A. C. Haza, and A. J. Mariano, "Submesoscale dispersion in the vicinity of the deepwater horizon spill," Proc. Natl. Acad. Sci. 111(35), 12693-12698 (2014).

<sup>40</sup>R. Lumpkin and M. Pazos, "Measuring surface currents with surface velocity program drifters: The instrument, its data, and some recent results," Lagrangian Analysis and Prediction of Coastal and Ocean Dynamics 39, 67 (2007).

<sup>41</sup>F. J. Beron-Vera and J. H. LaCasce, "Statistics of simulated and observed pair separations in the Gulf of Mexico," J. Phys. Oceanogr. 46, 2183-2199 (2016).

<sup>42</sup>A. Griffa, "Applications of stochastic particle models to oceanographic problems," in Stochastic Modelling in Physical Oceanography (Birkhäuser Boston, 1996), рр. 113–140."

<sup>43</sup>T. Rossby, M. Omand, J. Palter, and D. Hebert, "On rates of isopyc-nal dispersion at the submesoscale," Geophys. Res. Lett. **48**, e2021GL093526, https://doi.org/10.1029/2021GL093526 (2021).

44 M. Berta, A. Griffa, M. G. Magaldi, T. M. Özgökmen, A. C. Poje, A. C. Haza, and M. J. Olascoaga, "Improved surface velocity and trajectory estimates in the Gulf of Mexico from blended satellite altimetry and drifter data," J. Atmos. Oceanic Technol. 10, 1880-1901 (2015).

<sup>45</sup>N. Aksamit, T. Sapsis, and G. Haller, "Machine-learning mesoscale and submesoscale surface dynamics from Lagrangian ocean drifter trajectories," J. Phys. Oceanogr. 50, 1179–1196 (2020).

<sup>46</sup>G. Haller, N. O. Aksamit, and A. P. Encinas-Bartos, "Erratum: 'Quasiobjective coherent structure diagnostics from single trajectories' [Chaos 31, 043131 (2021)]," Chaos 32(5), 059901 (2022).

<sup>47</sup>R. Lumpkin, A. M. Treguier, and K. Speer, "Lagrangian eddy scales in the northern Atlantic Ocean," J. Phys. Oceanogr. 32(9), 2425-2440 (2002).

48 T. G. Shepherd, J. N. Koshyk, and K. Ngan, "On the nature of large-scale mixing in the stratosphere and mesosphere," J. Geophys. Res.: Atmos. 105(D10), 12433-12446, https://doi.org/10.1029/2000JD900133 (2000).

49 F. J. Beron-Vera, A. Hadjighasem, Q. Xia, M. J. Olascoaga, and G. Haller, "Coherent Lagrangian swirls among submesoscale motions," Proc. Natl. Acad. Sci. 116, 18251-18256 (2019).

<sup>50</sup>G. Haller, "Dynamically consistent rotation and stretch tensors from a dynamic polar decomposition," J. Mech. Phys. Solids **80**, 70–93 (2016). <sup>51</sup> A. Ruiz-Herrera, "Some examples related to the method of Lagrangian descrip-

tors," Chaos 25(6), 063112 (2015).

<sup>52</sup>A. Ruiz-Herrera, "Performance of Lagrangian descriptors and their variants in incompressible flows," Chaos 26(10), 103116 (2016).

<sup>53</sup>C. Mendoza, A. M. Mancho, and S. Wiggins, "Lagrangian descriptors and the assessment of the predictive capacity of oceanic data sets," Nonlinear Process Geophys. 21(3), 677-689 (2014).

<sup>54</sup>F. J. Beron-Vera and P. Miron, "A minimal Maxey-Riley model for the drift of Sargassum rafts," J. Fluid Mech. 904, A8 (2020).

<sup>55</sup>R. Lguensat, M. Sun, R. Fablet, P. Tandeo, E. Mason, and G. Chen, EddyNet: "A deep neural network for pixel-wise classification of oceanic eddies," in IGARSS 2018 IEEE International Geoscience and Remote Sensing Symposium (IEEE, 2018), рр. 1764–1767. 56 м. г.

<sup>6</sup>M. Farazmand and G. Haller, "Computing Lagrangian coherent structures from their variational theory," Chaos 22, 013128 (2012).

<sup>57</sup>O. R. Southwick, E. R. Johnson, and N. R. McDonald, "A simple model for sheddies: Ocean eddies formed from shed vorticity," J. Phys. Oceanogr. 46, 2961-2979 (2016).

58 O. R. Southwick, E. R. Johnson, and N. R. McDonald, "Potential vorticity dynamics of coastal outflows," J. Phys. Oceanogr. 47, 1021-1041 (2017).

<sup>59</sup>R. Lumpkin and G. C. Johnson, "Global ocean surface velocities from drifters: Mean, variance, El Niño-Southern Oscillation response, and seasonal cycle," J. Geophys. Res.: Oceans 118(6), 2992-3006, https://doi.org/10.1002/jgrc.20210 (2013).

<sup>60</sup>P. L. Richardson, "Eddy kinetic energy in the North Atlantic from surface drifters," J. Geophys. Res.: Oceans 88(C7), 4355-4367, https://doi.org/10.1029/ JC088iC07p04355 (1983).

<sup>61</sup>J. Martínez-Moreno, A. M. Hogg, A. E. Kiss, N. C. Constantinou, and A. K. Morrison, "Kinetic energy of eddy-like features from sea surface altimetry," J. Adv. Model. Earth Syst. 11(10), 3090-3105 (2019).

<sup>62</sup>D. Kang and E. N. Curchitser, "Gulf Stream eddy characteristics in a high-resolution ocean model," J. Geophys. Res.: Oceans 118, 4474-4487, https://doi.org/10.1002/jgrc.20318 (2013).

63 P. L. Richardson, A. E. Strong, and J. A. Knauss, "Gulf Stream eddies: Recent observations in the western Sargasso Sea," J. Phys. Oceanogr. 3, 297-301 (1973)

<sup>64</sup>C. E. Binding, T. A. Greenberg, and R. P. Bukata, "The MERIS maximum chlorophyll index; its merits and limitations for inland water algal bloom monitoring," J. Great Lakes Res. 39, 100-107 (2013).

<sup>65</sup>N. Tarshish, R. Abernathey, C. Zhang, C. O. Dufour, I. Frenger, and S. M. Griffies, "Identifying Lagrangian coherent vortices in a mesoscale ocean model," Ocean Modell, 130, 15-28 (2018).

<sup>66</sup>R. Lumpkin and L. Centurioni, "Global drifter program quality controlled 6-hour interpolated data from ocean surface drifting buoys," NOAA National Centers for Environmental Information, 2019.

<sup>67</sup>T. Özgökmen, GLAD experiment CODE-style drifter trajectories (low-pass filtered, 15 min interval records), northern Gulf of Mexico near DeSoto Canyon, July-October 2012. Distributed by: Gulf of Mexico Research Initiative Information and Data Cooperative (GRIIDC), Harte Research Institute, Texas A&M University-Corpus Christi (2013).

68 M. O. Williams, I. I. Rypina, and C. W. Rowley, "Identifying finite-time coherent sets from limited quantities of Lagrangian data," Chaos 25, 087408 (2015).